

Notes on Bundles on K3s

Eric Sharpe

Physics Department

Princeton University

Princeton, NJ 08544

ersharpe@puhep1.princeton.edu

In this technical note we describe a new (to the physics literature) construction of bundles on Calabi-Yaus. We primarily study this construction in the special case of K3 surfaces, for which interesting results can be obtained. For example, we use this construction to give plausibility arguments for a relationship between spaces of solutions of Hitchin's equations and moduli spaces of bundles on K3s. Also, in a recent paper it was proposed by C. Vafa that the mirror to a bundle on a Calabi-Yau n -fold is, in a particular sense, a supersymmetric n -cycle on the mirror Calabi-Yau. We use this new construction to observe that for the special case of K3s, Vafa's mirror data also specifies a bundle directly on the mirror K3, and so we potentially have a duality between bundles on any one K3 and other bundles on the mirror K3.

June 1998

1 Introduction

It was recently proposed by C. Vafa [1] that in an appropriate sense, the mirror to a Calabi-Yau n -fold with bundle should be a supersymmetric n -cycle on the mirror Calabi-Yau. More precisely, given a set of N Euclidean D- $(2n - 1)$ branes, the mirror should be thought of as a single D- $(n - 1)$ brane wrapped around a supersymmetric n cycle on the mirror Calabi-Yau. (The fact that we started with N branes is reflected in the homology class of the supersymmetric cycle on the mirror.)

In the special case of K3 surfaces, however, it is known that given a complex curve C in the K3 with a (reasonably well-behaved) line bundle on C , one can reconstruct a vector bundle on the entire K3 [2, 3, 4, 5]. Thus, in the special case of K3 surfaces, Vafa's construction potentially yields a duality between bundles on K3s and other bundles on mirror K3s.

More generally, we shall show how one can construct a bundle on a Calabi-Yau by specifying a complex codimension one subvariety C and a line bundle $\mathcal{L} \rightarrow C$. Essentially the same data has been used previously in constructions of bundles on elliptic Calabi-Yaus [17]; by contrast, our construction does not assume ellipticity. Although both constructions involve similar data, morally they take very different approaches to the problem of constructing bundles.

We shall also use the same construction to relate spaces of solutions of Hitchin's equations to moduli spaces of bundles on K3s, potentially opening the door to the application of integrable systems technology to understanding six-dimensional heterotic compactifications.

2 The construction

As is well-known, on a smooth variety one can define a line bundle by specifying a divisor. Loosely speaking, in this section we shall show how one can define higher-rank bundles given a divisor containing additional information.

Let C be a holomorphic curve in a K3, call it X . Let $\mathcal{L}$ be a line bundle on C . We shall need a technical restriction on $\mathcal{L}$ – it must be generated by its sections¹. Loosely speaking, for a line bundle to be generated by its sections means its global sections have the property that not all of them vanish at any point on C . More formally, this means we have a surjective map $H^0(\mathcal{L}) \otimes \mathcal{O}_C \rightarrow \mathcal{L}$.

In fact, we also have a surjective map $H^0(\mathcal{L}) \otimes \mathcal{O}_X \rightarrow \mathcal{L}'$, where $\mathcal{L}' = \iota_* \mathcal{L}$ and $\iota : C \hookrightarrow X$ is

¹For a bundle to be generated by its global sections means that for any prime ideal on the base, the images of the global sections generate the localization of the module [10, section II.5].

inclusion. (In other words, $\mathcal{L}'$ is a skyscraper² sheaf with support on C , such that $\mathcal{L}'|_C = \mathcal{L}$.) Denote the kernel of this map by $\mathcal{E}$, so we have the short exact sequence

$$0 \rightarrow \mathcal{E} \rightarrow H^0(C, \mathcal{L}) \otimes \mathcal{O}_X \rightarrow \mathcal{L}' \rightarrow 0 \quad (1)$$

The sheaf $\mathcal{E}$ is actually locally free, that is, $\mathcal{E}$ is a vector bundle on X , as opposed to a more general sheaf. This fact may seem somewhat surprising – since the short exact sequence (1) contains a skyscraper sheaf ($\mathcal{L}'$), one may be surprised that the kernel $\mathcal{E}$ is a vector bundle as opposed to a more general sheaf. However, the fact that the skyscraper sheaf has support at codimension one means the interpretation of $\mathcal{E}$ is slightly more subtle than one might at first have expected, and it turns out that $\mathcal{E}$ is actually a bundle. This phenomenon is commonly known in the mathematics literature as an elementary transformation, and these will be discussed in more detail (albeit in a totally different context) in [9]. For a more basic example of this phenomenon, let $\mathcal{I}$ be an ideal sheaf vanishing at codimension one, then we have the short exact sequence

$$0 \rightarrow \mathcal{I} \rightarrow \mathcal{O} \rightarrow \mathcal{O}/\mathcal{I} \rightarrow 0 \quad (2)$$

where $\mathcal{O}/\mathcal{I}$ denotes a skyscraper sheaf with support at codimension one (the vanishing locus of $\mathcal{I}$). Since $\mathcal{I}$ vanishes at codimension one, it is actually a line bundle, so the short exact sequence (2) is a prototype for the sequence (1).

In passing we should mention this trick hinges on the subvariety C having complex codimension one. One can certainly use this construction to create bundles on higher-dimensional Calabi-Yaus given a codimension one subvariety with a line bundle, however it cannot be applied if the subvariety has codimension greater than one. In particular, when Vafa's mirror construction is applied to Calabi-Yaus of higher dimension than K3, one does not recover a complex codimension one subvariety on the mirror, and so this construction does not apply. However, in special limits one might still be able to use a variant. For example, consider a four cycle in a Calabi-Yau four-fold. Suppose there exists a limit of complex structure in which the fourfold becomes reducible, with two components intersecting along a three-fold containing the four cycle. One could then use the construction above to create a bundle on the three-fold given a reasonably nice line bundle on the four-cycle, and then use a related construction to create a bundle on the four-fold from the one on the three-fold.

For reference, we shall repeat a few standard results. The bundle $\mathcal{E}$ has rank $h^0(C, \mathcal{L})$, $c_1(\mathcal{E}) = -[C]$, and $c_2(\mathcal{E}) = c_1(\mathcal{L})$. (As these Chern class computations are probably unfamiliar to the reader, they are reviewed in an appendix.) One can also compute

$$H^0(\mathcal{E}) = H^2(\mathcal{E}^\vee) = 0$$

² Actually, the term skyscraper sheaf is usually reserved for $\mathcal{O}_C$, in other words, when the restriction to the support is the structure sheaf (the trivial rank one bundle). More general sheaves with support on subvarieties of positive codimension are referred to as torsion sheaves. In this technical note we shall be sloppy on this point, blindly referring to all torsion sheaves as skyscraper sheaves.

$$\begin{aligned} H^1(\mathcal{E}) &= H^1(\mathcal{E}^\vee) = 0 \\ h^0(\mathcal{E}^\vee) &= h^0(\mathcal{L}) + h^1(\mathcal{L}) \end{aligned}$$

Note that the sequence (2) is not only a prototype for the sequence (1), but in fact is a simple example of the sequence (1). Suppose $\mathcal{L}$ is the trivial rank 1 bundle on C (the structure sheaf, more formally), then $\mathcal{L}$ is trivially generated by its sections and $h^0(C, \mathcal{L}) = 1$, so the corresponding bundle on X is a rank 1 bundle of $c_1 = -[C]$ – precisely the ideal sheaf $\mathcal{I}$ of sequence (2).

Given a holomorphic curve C and a reasonably nice line bundle $\mathcal{L} \rightarrow C$, we have shown how one may derive a bundle on the ambient K3. How does the moduli space of bundles of the same rank and Chern classes compare to the space of pairs $(C, \mathcal{L})$? The moduli space of bundles of rank r and Chern classes c_1, c_2 on K3 has dimension

$$\dim_{\mathbb{C}} \mathcal{M}_{K3} = 2rc_2 - (r-1)c_1^2 - 2(r^2 - 1)$$

so plugging in the values for the rank and Chern classes for a bundle derived from a pair $(C, \mathcal{L})$, and working in the special case that $h^0(C, \mathcal{L}^\vee \otimes K_C) = 0$ where K_C is the canonical bundle of C , we find that $\dim_{\mathbb{C}} \mathcal{M}_{K3} = 2g$, the same dimension as the space of pairs $(C, \mathcal{L})$ [11, 12].

3 Relation to Hitchin systems

A solution to Hitchin's equations [14, 15] is a pair $(\mathcal{E}, \phi)$ where $\mathcal{E}$ is a bundle on a Riemann surface C and $\phi \in H^0(C, \text{End}\mathcal{E} \otimes K_C) = H^1(C, \text{End}\mathcal{E})$, where K_C is the canonical bundle of C . This is the relevant worldvolume content of a set of D-branes wrapped on C [12]; ϕ is the Higgs scalar describing motions normal to C . Note that when we have a single D-brane, $\mathcal{E}$ is a line bundle, so $\text{End}\mathcal{E} = \text{Hom}(\mathcal{E}, \mathcal{E}) = \mathcal{E}^\vee \otimes \mathcal{E} = \mathcal{O}$, so $\phi \in \Gamma(K_C)$. (For a curve C inside a K3 (or any Calabi-Yau surface), a section of K_C specifies a deformation of C .)

Given that the dimensions of the moduli spaces match, it would appear quite plausible that moduli spaces of solutions of $U(1)$ Hitchin equations and certain moduli spaces of bundles on K3 are identical on some open subset of each, and that our construction of the previous section explicitly relates them. (Note that we have not proven this – we are merely observing that under the circumstances it seems natural.) As a check, note that it is known that both are typically birational to a Hilbert scheme of points on K3 [12, 13]. (Related results have also been obtained in [6, 7, 8].) This result seems to be special to K3 surfaces – it does not seem to hold for curves in other surfaces.

Assuming that moduli spaces of bundles on K3 and moduli spaces of pairs $(C, \mathcal{L})$ are related, there should exist a description of Mumford-Takemoto and Gieseker stability of

bundles on K3 in terms of pairs $(C, \mathcal{L})$. Unfortunately, we have not yet been able to see the corresponding versions of these stability conditions.

Clearly not every bundle on a K3 can be viewed as descending from a holomorphic curve C with line bundle $\mathcal{L} \rightarrow C$, if for no other reason than the fact that not every bundle on K3 will satisfy $H^0(\mathcal{E}) = H^1(\mathcal{E}) = 0$. Also, the rank and Chern classes of bundles descending from pairs $(C, \mathcal{L})$ are not independent, but are related by Riemann-Roch, so again not all bundles can be described in this fashion.

Although not every bundle on a K3 can be constructed directly in this fashion, it should still be possible to describe many of them – if $\mathcal{E}$ denotes a bundle constructed as in equation (1), then by tensoring $\mathcal{E}$ and $\mathcal{E}^\vee$ with line bundles it is possible to recover many more bundles.

So far we have argued that, as our construction builds bundles on K3s from pairs $(C, \mathcal{L})$, it implicitly relates solutions of $U(1)$ Hitchin equations to bundles on K3. In the next section we shall speak to Hitchin systems with larger gauge groups, and argue that essentially nothing new happens.

4 Higher rank bundles on a curve

So far we have only spoken about constructing bundles on a K3 given a rank 1 bundle on a curve. What if instead we have a higher rank bundle on the same curve? Can one construct additional bundles on the K3, not obtainable with mere line bundles?

The answer seems to be no, by studying more general vector bundles on curves we do not learn anything new. Physically this should be clear: a rank m vector bundle on a curve C describes a collection of m D-branes wrapped on C , which should be physically equivalent to a single D-brane wrapped on a curve in the homology class of mC .

Mathematically this intuition also works out. Let us work through this in nontrivial detail, so as to be more convincing. Let $\mathcal{F} \rightarrow C$ denote a rank m vector bundle on C , generated by its sections. Define a bundle $\mathcal{E}$ on the ambient K3, denoted X , as the kernel

$$0 \rightarrow \mathcal{E} \rightarrow H^0(C, \mathcal{F}) \otimes \mathcal{O}_X \rightarrow \mathcal{F}' \rightarrow 0$$

where $\mathcal{F}' = \iota_* \mathcal{F}$. It is straightforward to compute

$$\begin{aligned} c_1(\mathcal{F}') &= mC \\ c_2(\mathcal{F}') &= -c_1(\mathcal{F}) + \frac{1}{2}m(m+1)C^2 \end{aligned}$$

and from this we find

$$c_1(\mathcal{E}) = -mC$$

$$c_2(\mathcal{E}) = c_1(\mathcal{F}) + m(m-1)(g-1)$$

where g is the genus of C .

The reader should now (correctly) guess that a vector bundle $\mathcal{F} \rightarrow C$ describes the same $\mathcal{E} \rightarrow X$ as a line bundle $\mathcal{L} \rightarrow C'$ where C' is in the homology class of mC (so C' has genus $g' = m^2(g-1) + 1$), and

$$c_1(\mathcal{L}) = c_1(\mathcal{F}) + m(m-1)(g-1)$$

As a check, in the case that $h^0(C, \mathcal{F}^\vee \otimes K_C) = 0$, it is easy to compute that the dimension of the moduli space of bundles on K3 of rank $h^0(C, \mathcal{F})$, $c_1 = -mC$, and $c_2 = c_1(\mathcal{F}) + m(m-1)(g-1)$ is precisely $2g'$. We have been slightly sloppy; the precise curve $C' \in [mC]$ is specified by a choice of section $\phi \in \Gamma(C, \text{End}\mathcal{F} \otimes K_C)$, corresponding to the Higgs scalar of the wrapped D-brane worldvolume describing motions normal to C .

In fact, such a relation³ between vector bundles on curves (paired with sections of $\text{End}\mathcal{F} \otimes K_C$) and line bundles on coverings is known to exist – see for example [14, section 5] and [16]. We shall not repeat their explanation here.

To summarize, given a vector bundle $\mathcal{F}$ on a curve C in some K3, together with a section $\phi \in \Gamma(\text{End}\mathcal{F} \otimes K_C)$, we can construct a covering space C' of C inside the total space of the normal bundle to C , and a line bundle $\mathcal{L} \rightarrow C'$. Thus, the data specifying a solution of Hitchin's equations, for any gauge group, on a curve in a K3 will specify a bundle on the entire K3. Moreover, the data defining a rank n bundle on a curve C can be duplicated by a line bundle on a cover $C' \rightarrow C$.

5 Conclusions

In this paper we have described a new (to the physics literature) construction of bundles on Calabi-Yaus, in which bundles are specified by a complex codimension one subvariety C and a line bundle $\mathcal{L} \rightarrow C$. We have applied this construction to the special case of K3 surfaces, and in particular noted a connection between moduli spaces of bundles on K3s and moduli spaces of solutions of Hitchin's equations.

³We are being slightly sloppy. The references cited above describe a connection between vector bundles (and sections) on some curve and line bundles on a ramified cover. However coincident D-branes are not precisely described by a single D-brane on a cover (as elements of the cover are generically separated by a finite distance), though the two situations are certainly in the same homology class. Coincident D-branes are best described scheme-theoretically, and in this instance would be described by a thick curve. It seems likely that there is an identical correspondence between vector bundles on curves and line bundles on thick curves, but to our knowledge the details of such a correspondence have not been worked out in the mathematics literature.

We should point out that we have been rather sloppy about the physics of this construction. We have outlined how one can construct a bundle $\mathcal{E}$ from a holomorphic curve C and a line bundle $\mathcal{L} \rightarrow C$, but it is not clear that $\mathcal{E}$ should be physically preferred over $\mathcal{E}^\vee$ or $\mathcal{E} \otimes \mathcal{O}(D)$ for some line bundle $\mathcal{O}(D)$ on K3. More precisely, although we have a D-brane wrapping C , we do not have a D-brane wrapping the K3 – our construction of a bundle is purely formal in nature.

The reader might wonder if one could use our extension of Vafa’s duality [1] to create a $\mathbf{Z}_2$ symmetry on the space of bundles on K3s, assuming we found a way to eliminate the ambiguity mentioned in the paragraph above. In particular, given any bundle on a K3, we can use Vafa’s prescription to associate a D-brane wrapped on a curve in the mirror K3, then associate a bundle as above and repeat. However we should point out that the bundle we create is not clearly associated to any D-branes wrapped on the mirror K3, so there is no physics reason to believe one could get a duality symmetry. Moreover, if one applies this construction twice (pretending that the bundle on the mirror K3 lives on some “virtual” brane), in general one will not recover the original bundle on the original K3 – one will recover a bundle descending from a pair $(C, \mathcal{L})$, but the original bundle need not have been associated to such a pair.

Nevertheless, it might be possible to find a $\mathbf{Z}_2$ symmetry on the subspace of bundles which happen to be associated to pairs $(C, \mathcal{L})$, although there is no purely physics argument to reach such a conclusion. The ambiguity in which bundle to associate to a pair $(C, \mathcal{L})$ might be fixed by the requirement of obtaining a $\mathbf{Z}_2$ symmetry. Unfortunately, we have not yet been able to reach any definite conclusions in this matter. If such a $\mathbf{Z}_2$ symmetry exists, it may or may not amount to a special case of a Fourier-Mukai transform – again, we do not have anything definite to say in the matter.

The reader may wonder if there is any connection between the work described in this technical note and the description of bundles on elliptic K3s with section given in [17]. In both instances bundles are described in terms of some curve in a K3 with a line bundle on the curve. However the similarity between the two approaches ends there – morally our approach and theirs are quite different. If nothing else, our approach does not require that the K3 be elliptic (merely algebraic).

In this technical note we only spoke to the description of $GL(n, \mathbf{C})$ bundles on K3; what about bundles with other gauge groups? In principle, one could attempt to use the same construction to build $GL(n, \mathbf{C})$ bundles with reducible structure group. Work on this matter is in progress.

Finally, note that one might also be able to use this construction to find four-dimensional analogues of many two-dimensional quantities. For example, it might conceivably be possible to construct a four-dimensional version of the Verlinde formula directly from its usual two-dimensional form. (Four-dimensional analogues of the Verlinde formula have been previously

discussed in [20, 21].)

6 Acknowledgements

We would like to thank P. Aspinwall, M. Bershadsky, A. Mikhailov, D. Morrison, and E. Witten for useful conversations, and especially T. Gomez for extremely close readings of rough drafts.

A Chern class computations

The reader may wonder how to compute the Chern classes of the skyscraper sheaf $\mathcal{L}'$ appearing in equation (1), as well as the higher-rank skyscraper sheaves appearing later. We shall begin this appendix with a relatively intuitive (though not completely rigorous) derivation of the Chern classes of these sheaves, then shall outline how to derive the results rigorously with Grothendieck-Riemann-Roch.

We shall begin with an intuitive derivation of the Chern classes. The skyscraper sheaf $\mathcal{L}' \rightarrow X$ appearing in equation (1) can often⁴ be obtained by restriction of a line bundle $\mathcal{O}(D) \rightarrow X$ to the curve C : $\mathcal{L}' = \mathcal{O}_C \otimes \mathcal{O}(D)$ for some D , with $c_1(\mathcal{L}) = C \cdot D$. Given the resolution of the skyscraper sheaf

$$0 \rightarrow \mathcal{O}_X(-C) \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_C \rightarrow 0$$

we find the resolution

$$0 \rightarrow \mathcal{O}_X(-C + D) \rightarrow \mathcal{O}_X(D) \rightarrow \mathcal{L}' \rightarrow 0$$

and so $c_1(\mathcal{L}') = C$, $c_2(\mathcal{L}') = C^2 - C \cdot D$. Given the Chern classes of $\mathcal{L}$, the Chern classes of $\mathcal{E}$ are of course obtained as $c(\mathcal{E}) = c(\mathcal{L}')^{-1}$. Chern classes of higher-rank skyscraper sheaves can be derived with the splitting principle and the same general methods.

Although the derivation above is relatively intuitive, it is not always rigorous, simply because $\mathcal{L}'$ is not always the tensor product of a line bundle on X with $\mathcal{O}_C$. To recover the Chern classes in complete generality, we shall use a specialization of the Grothendieck-Riemann-Roch theorem [10, section A.5], which in these circumstances states (see [18, section 9.1] or [19, section 15.1])

$$ch(\iota_* \mathcal{F}) = \iota_* (ch(\mathcal{F}) \cdot td(\mathcal{N})^{-1})$$

⁴It is not always the case that $\mathcal{L}'$ is the restriction of a line bundle on X , however this assumption leads to a simple derivation. In the event that $\mathcal{L}'$ is the restriction of a line bundle, this derivation is rigorous.

where $\mathcal{F}$ is a coherent sheaf on C , and $\mathcal{N}$ is the normal bundle to C in X . (In fact, $\mathcal{N} = K_C$, the canonical bundle of C .)

Now, ι_* acts in the obvious way on elements of the Chow ring, so $\iota_*(1) = C$, $\iota_*c_1(\mathcal{L}) = c_1(\mathcal{L})$ (where the right side is now interpreted as defining an element of $H^4(X)$), and [10, section A.3] $\iota_*c_1(K_C) = C^2$ because K_C is the normal bundle to C in X .

With a small amount of work, we finally recover

$$\begin{aligned} c_1(\iota_*\mathcal{L}) &= \iota_*(1) \\ &= C \\ \frac{1}{2}c_1(\iota_*\mathcal{L})^2 - c_2(\iota_*\mathcal{L}) &= \iota_*\left(c_1(\mathcal{L}) - \frac{1}{2}c_1(K_C)\right) \\ &= c_1(\mathcal{L}) - \frac{1}{2}C^2 \end{aligned}$$

which is the desired result.

References

- [1] C. Vafa, “Extending Mirror Conjecture to Calabi-Yau with Bundles,” [hep-th/9804131](#).
- [2] R. Lazarsfeld, “Brill-Noether-Petri without Degenerations,” *J. Diff. Geom.* **23** (1986) 299.
- [3] A. N. Tyurin, “Cycles, Curves and Vector Bundles on an Algebraic Surface,” *Duke Math. J.* **54** (1987) 1.
- [4] M. Green and R. Lazarsfeld, “Special Divisors on Curves on a K3 Surface,” *Inv. Math.* **89** (1987) 357.
- [5] T. Gomez, “Brill-Noether Theory on Singular Curves and Vector Bundles on K3 Surfaces,” Ph.D. thesis, [alg-geom/9710029](#).
- [6] R. Donagi, L. Ein, and R. Lazarsfeld, “Nilpotent Cones and Sheaves on K3 Surfaces,” [alg-geom/9504017](#), in *Birational Algebraic Geometry*, ed. Y. Kawamata and V. V. Shokurov, Contemporary Mathematics 207, American Mathematical Society, 1997.
- [7] R. Donagi and E. Markman, “Spectral Covers, Algebraically Completely Integrable Hamiltonian Systems, and Moduli of Bundles,” [alg-geom/9507017](#), in *Integrable Systems and Quantum Groups*, Lecture Notes in Math 1620, Springer, Berlin, 1996.
- [8] S. Mukai, “Symplectic Structure of the Moduli Space of Sheaves on an Abelian or K3 Surface,” *Invent. Math.* **77** (1984) 101.

- [9] E. Sharpe, in preparation.
- [10] R. Hartshorne, *Algebraic Geometry*, Springer-Verlag, New York, 1977.
- [11] A. Strominger, S.-T. Yau, and E. Zaslow, “Mirror Symmetry is T-Duality,” Nucl. Phys. **B479** (1996) 243, [hep-th/9606040](#).
- [12] M. Bershadsky, V. Sadov, and C. Vafa, “D-Branes and Topological Field Theories,” Nucl. Phys. **B463** (1996) 420, [hep-th/9511222](#).
- [13] C. Vafa, “Instantons on D-Branes,” Nucl. Phys. **B463** (1996) 435, [hep-th/9512078](#).
- [14] N. Hitchin, “Stable Bundles and Integrable Systems,” Duke Math. J. **54** (1987) 91.
- [15] N. Hitchin, “The Self-Duality Equations on a Riemann Surface,” Proc. London Math. Soc. (3) **55** (1987) 59.
- [16] A. Beauville, M. S. Narasimhan, and S. Ramanan, “Spectral Curves and the Generalized Theta Divisor,” J. reine angew. Math. **398** (1989) 169.
- [17] R. Friedman, J. Morgan, and E. Witten, “Vector Bundles and F Theory,” Comm. Math. Phys. **187** (1997) 679, [hep-th/9701162](#).
- [18] W. Fulton, *Introduction to Intersection Theory in Algebraic Geometry*, Regional Conference Series in Mathematics Number 54, American Mathematical Society, Providence, Rhode Island, 1984.
- [19] W. Fulton, *Intersection Theory*, Springer-Verlag, Berlin, 1984.
- [20] N. Nekrasov, “Four Dimensional Holomorphic Theories,” Princeton University Ph.D. thesis, unpublished.
- [21] A. Losev, G. Moore, N. Nekrasov, and S. Shatashvili, “Four-Dimensional Avatars of Two-Dimensional RCFT,” Nucl. Phys. Proc. Suppl. **46** (1996) 130, [hep-th/9509151](#).